

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number:

0 337 601
A1

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 89301924.0

(51) Int. Cl. 4: **C07D 487/22 , A61K 31/40 ,**
///(C07D487/22,257:00,209:00)

(22) Date of filing: 27.02.89

(30) Priority: 11.03.88 GB 8805849

(43) Date of publication of application:
18.10.89 Bulletin 89/42

(24) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

(71) Applicant: **EFAMOL HOLDINGS PLC**
Efamol House Woodbridge Meadows
Guildford Surrey GU1 1BA(GB)

(72) Inventor: **Bonnett, Raymond c/o Queen Mary**


College (QMC)
Mile End Road
London E1 4NS(GB)
Inventor: **Berenbaum Morris Cyril**
c/o St. Mary's Hospital Medical School
London W2 1PG(GB)

(74) Representative: **Caro, William Egerton et al**
J. MILLER & CO. Lincoln House 296-302 High
Holborn
London WC1V 7JH(GB)

(54) **Porphyrins and cancer treatment.**

(57) This invention provides novel compounds for therapy of tumours susceptible to necrosis when an appropriate compound is administered followed by illumination with light of a wavelength absorbed by the compound, the compounds being dihydro porphyrins of the formula, or the corresponding tetra-hydro porphyrins:

EP 0 337 601 A1



where $n = 1$ to 3 and each substituent R , the same or different and at the same or different position in its respective substituent phenyl ring, is an hydroxy group.

PORPHYRINS AND CANCER TREATMENT

FIELD OF THE INVENTION

This invention relates to certain porphyrins and their use in cancer treatment.

5

BACKGROUND

Haematoporphyrin derivatives of uncertain specific composition (HpD) have been used in cancer treatment, having been found to locate in tumours and other tissues after injection into the bloodstream, and to sensitise cells to light irradiation (transport in the blood is believed to be largely in association with the serum albumin). Irradiation, with a laser or other source, may be direct or indirect, for example using fibre optics. Irradiated cells, unless deeply pigmented, are rapidly killed to a depth depending on the light penetration. The mechanism of cell killing in phototherapy is believed to be largely by production of singlet oxygen. This is produced by transfer of energy from the light-excited porphyrin molecule to an oxygen molecule. Singlet oxygen is highly reactive. It is believed to oxidise cell membranes so that they are damaged and become incapable of exerting their function of controlling the cell's internal operation. This rapidly leads to cell death.


Selectivity of damage for various tissues, including tumours, is governed by a variety of factors, including selective uptake, selective vulnerability and selective ability to repair damage (cf. Albert A.A. "Selective Toxicity" fifth edition, Chapman & Hall, London 1973). In tumours an additional factor is the precarious nature of tumour blood supply and its particular vulnerability to photo-dynamic damage as compared with the robust and well-established blood supply of normal tissues. All the above factors vary from one tissue to another and from one compound to another, so that the pattern of tumour and normal tissue damage may be expected to vary.

In EP-A-0,186,962, the present applicants made proposals to improve on HpD by finding well characterised and thus more exactly controllable compounds. Other aims were to find compounds activated by light at wavelengths longer than those used to activate HpD so as to exploit deeper penetration of longer wavelength radiation, and to increase effectiveness generally, as in many anatomical sites, such as the brain, HpD has been found to sensitise normal cells unduly as well as cancerous cells. The wavelength at which a photo-sensitising compound is activated is one factor in its *in vivo* effectiveness. Other things being equal, the longer the activating wavelength within the visible range, the greater the tissue penetration of light and therefore the greater the depth of damage. Thus, compounds activated at 650-660nm might be expected to produce greater depths of damage than HpD, which is activated at 625-630nm. Successful photo-therapy depends on the ability to produce severe tumour damage without unacceptable damage to contiguous normal tissues. The compounds used according to EP-A-0,186,962 are porphyrins tetra-meso-substituted by aromatic groups bearing hydroxy, amino or sulphhydryl groups, some new as compounds others old.


The present invention also concerns medicaments for therapy of tumours susceptible to necrosis when an appropriate compound is administered to locate in the tumour followed by illumination of the tumour with light of a wavelength absorbed by the compound, but the compounds are now and the invention therefore lies primarily in them. They are dihydro porphyrins (chlorins) I, and corresponding tetra-hydro porphyrins II and III, of the formulae:

45

50



wherein each Ar, the same or different, is an aromatic group with one or more hydroxy (-OH) substituent groups. For example, and showing the ring numbering, the dihydro porphyrins (and the tetra-hydro porphyrins correspondingly) may be:-



wherein each R (one or more in each ring $n = 1$ to 3) is an ortho, meta or para positioned hydroxy substituent group, particularly to give polyhydroxyphenyl compounds. Said substituent groups may be in the same or different positions on their respective aromatic groups. Any said substituent group may be free or itself substituted, for example by alkyl or acyl groups preferably C_1 to C_4 and the compounds when in such form are within the claims herein. The nucleus or the substituent rings may be substituted further, provided pharmacological tolerability, appreciable solubility in water (required so that the drug may be administered intravenously to ensure rapid distribution to the tumour), absorption of light at the red end of the spectrum, and take up in cancerous tissue are retained, and again the compounds when in such form are to be understood as within the claims herein.







Any of the compounds further may be in the form of derivatives such as salts at acidic or basic centres, metal complexes (e.g. Zn, Ga), or hydrates or other solvates particularly with lower, e.g. C_1 - C_4 aliphatic alcohols and again such derivatives are within the claims.

It is preferred that one or more of the said substituent groups should be of a kind and in a form able to ionise at physiological pH, to increase the absorption in the red part of the spectrum, that is in the portion that most effectively penetrates tissue. Compounds not very soluble in themselves may be solubilised by the presence of suitable groups such as sulphonate groups.

The invention further extends to the medicaments, to their preparation and to the method of therapy itself.

Various specific compounds within the claims are given in Table 1 below:-

TABLE 1

NUMBER	NATURE OF "Ar" IN FORMULA I or II	CODE NAME (DIHYDRO or TETRA-HYDRO FORM OF)
1		HK7
2		acetyl HK7
3		ortho-HK7
4		ortho-HK7 methyl ether
5		meta-HK7
6		acetyl meta-HK7

The invention is illustrated in the following Examples, in which the compounds are made by the reduction of the corresponding porphyrins, without protection of the latter. The general method of reduction is not new (Whitlock et al., J. Amer. Chem. Soc., 1969, 91, 7485) but the products are new. It is unexpected that they can be obtained without prior protection of phenolic hydroxy functions, especially when an oxidative step is required in the work up. When required, a chloranil dehydrogenation step is included to oxidise tetra-hydro porphyrins formed back to the dihydro form.

10 EXAMPLE 1

Synthesis of 2,3-dihydro-5,10,15,20-tetra(p-hydroxyphenyl) porphyrin ("Dihydro HK7")

15 5,10,15,20-Tetra(p-hydroxyphenyl)porphyrin (163mg, 0.24 mmoles), p-toluenesulphonhydrazide (90mg, 0.48 mmoles), anhydrous potassium carbonate (300mg) and anhydrous pyridine (11.25 ml) were stirred with nitrogen flushing for 20 minutes. The mixture was heated (100-105 °C) for 6.5 hours under nitrogen: further quantities of p-toluenesulphonhydrazide (97mg in 0.3 ml anhydrous pyridine on each occasion) were added after 2 hours and 4 hours. The reaction mixture was treated with ethyl acetate (75 ml) and distilled water (37.5 ml) and digested on a steam bath for 1 hour. The organic layer was separated and washed in turn with hydrochloric acid (2M, 75 ml), distilled water (75 ml) and saturated sodium bicarbonate solution (75 ml).

o-Chloranil (total 166mg, 0.675 mmoles) was added in portions to the stirred organic solution at room temperature until the absorption peak at 735 nm had disappeared.

25 The mixture was washed with aqueous sodium bisulphite (5%, 2 x 75 ml), distilled water (75 ml), sodium hydroxide (0.01M, 100 ml) and saturated sodium bicarbonate (80 ml) and dried over anhydrous sodium sulphate. The filtered solution was evaporated to dryness and the residue was crystallised from methanol-distilled water to give 77 mg (middle cut, 47%) of 2,3-dihydro-5,10,15,20-tetra(p-hydroxyphenyl)-porphyrin as a purple solid. Mass spectrum (FAB) Found (M + H)⁺ 681.253 C₄₄H₃₂N₄O₈ requires M + 1 = 681.250. Nmr spectrum in accord with structure. Electronic spectrum λ_{max} (MeOH) (ε) 285 (17000), 295 (16900), 418 (143000), 520 (10000), 550 (9000), 597 (5100), and 651 (18600). R_f 0.49 on Merck silica gel irrigated with MeOH:CHCl₃ (1:4).

35 EXAMPLE 2

Synthesis of 2,3-dihydro-5,10,15,20-tetra(m-hydroxyphenyl) porphyrin ("Dihydro meta-HK7")

40 The method was as described above starting with the following reagents: 5,10,15,20-tetra(m-hydroxyphenyl) porphyrin (160mg, 0.23 mmoles), p-toluenesulphonhydrazide (90mg, 0.48 mmoles), anhydrous potassium carbonate (303mg), anhydrous pyridine (11.25 ml), and o-chloranil (106mg, 0.43 mmoles).

The middle cut from the crystallisation from methanol-distilled water gave 59mg (37%) of 2,3-dihydro-5,10,15,20-tetra(m-hydroxyphenyl)porphyrin as a purple solid. Mass spectrum (FAB) (M + H)⁺ 681. C₄₄H₃₂N₄O₄ requires M 680. Electronic spectrum λ_{max} (MeOH) (ε) 284 (16900), 306 (15600), 415 (146000), 516 (11000) 543 (7300), - 591 (4400), and 650 (22400). R_f 0.52 on Merck silica gel irrigated with MeOH:CHCl₃ (1:4).

50 EXAMPLE 3

Synthesis of 2,3-dihydro-5,10,15,20-tetra (o-hydroxyphenyl) porphyrin ("Dihydro ortho-HK7").

55 The method was as described above starting with the following reagents: 5,10,15,20-tetra (o-hydroxyphenyl) porphyrin (163 mg, 0.24 m moles), anhydrous potassium carbonate (303 mg), anhydrous pyridine (11.25 ml), and o-chloranil (132 mg, 0.54 m moles).

Crystallisation from methanol-distilled water gave 105 mg (64%) of 2,3-dihydro-5,10,15,20 -tetra(o-

hydroxyphenyl) porphyrin as a dark purple crystalline solid. It was a mixture of atropisomers (R_f 0.23, 0.29, 0.38 and 0.43 on silica gel irrigated with 1% MeOH in CHCl_3). Mass spectrum (FAB) ($M+H$)⁺ 681. $\text{C}_{44}\text{H}_{32}\text{N}_4\text{O}_4$ requires M 680. Electronic spectrum λ_{max} (MeOH) (ϵ) 415 (90700), 515 (8400), 542 (5600), 597 (3800), and 651 (16000).

5

EXAMPLE 4

10 Synthesis of 7,8,17,18-tetrahydro-5,10,15,20-tetra(m-hydroxyphenyl) porphyrin

5,10,15,20-tetra(m-hydroxyphenyl)porphyrin (109 mg, 0.16 mmoles), p-toluenesulphonylhydrazide (60 mg), anhydrous potassium carbonate (200 mg) and anhydrous pyridine (7.5 ml) were stirred with nitrogen flushing for 20 min. The mixture was heated (100° - 105°C) for a total of 12 hours under nitrogen, further
 15 quantities of p-toluenesulphonylhydrazide being added every 1.5 hour (60 mg in 0.2 ml pyridine on each occasion). The reaction mixture was treated with ethyl acetate (100 ml) and water 50 ml) and digested on a steam bath for 1 hour. The organic layer was separated, and washed with hydrochloric acid (2M, 50 ml), and then with phosphoric acid (56%, 4 x 50 ml, to remove chlorin). The organic layer was washed in turn with distilled water (50 ml), sodium bicarbonate (50 ml), distilled water (50 ml), hydrochloric acid (2M,
 20 50ml), and distilled water (50 ml), and dried over anhydrous potassium carbonate. After filtration and removal of solvent the residue was crystallised from methanol-water to give the title compound as a green solid (29 mg, 27%). R_f 0.47 on silica gel irrigated with MeOH- CHCl_3 (1:4).

λ (MeOH, nm) (ϵ) 352 (92,000), 361 (114,000), 372 (129,000), 516 (50,000) and 735 (91,000).

δ (d_6 -DMSO) (250 MHz):- 9.69 (s, 3.8H, OH); 7.95 (s, 4.OH, β -pyrrole H), 7.47 (t, 5.2H, ArH); 7.21 (d,
 25 4.2H, ArH); 7.18 (s, 4.2H, ArH); 7.04 (d, 4.OH, ArH); 3.95 (s, 7.5H, CH_2); and -1.54 (s, 2.OH, NH).

The activity of the compounds has been demonstrated in tests given in Table 2 below, the dihydro compounds being referred to as chlorins and the tetra-hydro compounds as bacteriochlorins.

TABLE 2

30

Tumour photo-necrosis with chlorins (C) and a bacteriochlorin (BC) of the meso -tetra(hydroxyphenyl) (THP)			
Photosensitiser	Dose $\mu\text{M/Kg}$	Wavelength* (nm)	Depth of Tumour Necrosis mm \pm SE**
p-THPC	6.25	653	3.50 \pm 0.54 (10)
	3.125	653	2.13 \pm 0.50 (10)
m-THPC	0.75	652	5.41 \pm 0.39 (19)
	0.375	652	3.79 \pm 0.28 (6)
o-THPC	6.25	652	4.16 \pm 0.27 (14)
	3.125	652	3.69 \pm 0.74 (9)
m-THPBC	0.39	741	5.22 \pm 1.21 (8)

* The total energy administered was 10 J cm^{-2} throughout.

45

** Number of tumours in parenthesis.

The method of testing is that the PC6 plasma cell tumour, obtained initially from the Chester Beatty Research Institute, was grown by inoculating $0.3 - 0.6 \times 10^6$ cells subcutaneously in female BALB/c mice. It was used about two weeks later, when it was 12 - 13 mm in its longest diameter and 6 - 7 mm deep.
 50 Sensitisers were injected in dimethyl sulphoxide intra-peritoneally (2.5 ul g^{-1}) and, twenty four hours later, skin over the tumours was depilated, the mice anaesthetised and tumours exposed to 10 J cm^{-2} light. Wavelengths for illumination were the longest wavelength absorption peaks in the red in solutions of sensitiser in fetal calf serum, and were 652 - 653 nm for the compounds of Examples 1 to 3 and 741 nm for the compound of Example 4. The light source was a copper vapour laser with an output of 10 to 12 watts (Cu 10 laser, Oxford Lasers Ltd) pumping a D2 10K dye laser (Oxford Lasers Ltd). The dye was rhodamine 640 (Applied Photophysics Ltd) for illumination at 652 - 653 nm and LDS-722 for illumination at 741 nm. Light intensity at the tumour surface was kept below 0.3 W cm^{-2} , where thermal effects were undetectable.

55

Twenty four hours after illumination at 652 - 653 nm and LDS 722 for illumination at 741 nm, 0.2 ml 1%

Evans blue (Sigma) in saline was given intravenously and tumours were removed one hour later and fixed in formol-saline. The fixed tumours were sliced at right angles to the surface and the depth of necrosis measured with a dissecting microscope fitted with an eye piece graticule.

In preliminary tests the new compounds have shown similar results as regards skin sensitivity to those given in EP-A-0,186,962. Moreover, the para and meta dihydroxy compounds have been shown specifically, improved properties in other respects which may be expected in all the compounds. The results of a number of experiments are given in Tables 3 and 4 below.

TABLE 3

DEGREE OF TUMOUR NECROSIS OBTAINED AT AN EQUIVALENT MUSCLE DAMAGE LEVEL		
Muscle Oedema	Tumour Necrosis (mm)	
(% increase in muscle weight)	Corresponding Compounds of EP-A-0186962	Compounds of Examples 1 and 2 above
	Para	Para
10%	2.25	3.5
20%	3.0	4.0 approx.
	Meta	Meta
10%	3.5	5.5-6
20%	4.0	6.5

TABLE 4

COST IN TERMS OF MUSCLE DAMAGE FOR SPECIFIED DEPTHS OF TUMOUR NECROSIS		
Depth of Tumour Necrosis (mm)	Muscle Oedema (% Increase in Weight)	
	Corresponding Compounds of EP-A-0186962	Compounds of Examples 1 and 2 above
	Para	Para
3.0	20	8
3.5	22	10
	Meta	Meta
3.5	10	2
4.0	20	3
5.0	36	5
5.5	49	7

CONCLUSION

Tables 3 and 4 show that, for specified levels of muscle damage, the compounds of Examples 1 and 2 produce more tumour necrosis than the corresponding compounds of EP-A-0,186,962 and, for specified levels of tumour necrosis, cause considerably less muscle damage than the corresponding compounds of EP-A-0,186,962.

This type of selectivity is particularly important in the photo-therapy of tumours in anatomical sites where muscle is a major constituent, e.g., in cancer of the head and neck, and in cancer of the large intestine and rectum with possible involvement of abdominal wall or pelvis.

In administration the compounds, as with known tumour sensitisers, will be given intravenously in a blood compatible preparation. Conveniently an aqueous solution is used but if water solubility is limited, a solution in an organic, non-toxic polar solvent such as ethanol or a mixture of such solvents may for example be prepared. A further route to water solubility in the sense of transport in the aqueous environment of the bloodstream is the preparation of liposomes to be administered in aqueous media. Therapy according to the invention is thus administered in suitable form in an amount determined by the general toxicity of the particular compound and its level of necrotic effect, allowing tumour location to take place, and illuminating with light of a wavelength reaching the tumour and exciting the compound, at a dose effecting tumour necrosis without unacceptably damaging normal tissue. All this is in per se known manner, not requiring detailed discussion on its optimisation.

15


Claims

1. Compounds being pharmacologically acceptable tumour locating dihydro or tetra-hydro porphyrins of the formulae below; methods of making medicaments for therapy of tumours susceptible to necrosis when such a compound is administered to locate in the tumour followed by illumination of the tumour with light of a wavelength absorbed by the compound, said methods making use of the compounds by associating them with a pharmaceutical diluent or carrier; and such medicaments and therapy as such; the compounds being of the formulae:-

25

30


35



40

45


50



wherein each Ar, the same or different, is an aromatic group with one or more hydroxy (-OH) substituent groups.


55

2. The subject matter of claim 1, the compounds being of the formula:-



or the corresponding tetra-hydro porphyrins, where $n = 1$ to 3 and each substituent R , the same or different and at the same or different position in its respective substituent phenyl ring, is an hydroxy group.

3. The subject matter of claim 2 where each substituent



is an *o*-hydroxyphenyl, *m*-hydroxyphenyl or *p*-hydroxyphenyl group.

4. The subject matter of claim 1, 2 or 3 wherein each hydroxy group is free rather than in a substituted form.

5. The subject matter of claims 1, 2, 3 or 4 the compounds being synthesis of 2,3-dihydro-5,10,15,20-tetra(*p*-hydroxy-phenyl) porphyrin; synthesis of 2,3-dihydro-5,10,15,20-tetra(*m*-hydroxy-phenyl) porphyrin; synthesis of 2,3-dihydro-5,10,15,20-tetra (*o*-hydroxyphenyl) porphyrin and synthesis of 7,8,17,18-tetrahydro-5,10,15,20-tetra(*m*-hydroxyphenyl) porphyrin.



European Patent
Office

PARTIAL EUROPEAN SEARCH REPORT
which under Rule 45 of the European Patent Convention
shall be considered, for the purposes of subsequent
proceedings, as the European search report

Application number
EP 89301924.0
- page 1 -

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 7)
D, A	EP - A - 0186962 (EFAMOL LTD) * claims *	1-4	C07D487/22 A61K31/40 //(C07D487/22, 257:00,209:00, 209:00,209:00, 209:00)
A	--- JOURNAL OF THE AMERICAN CHEMICAL SOCIETY, vol. 95, no. 17, 22.08.73, pages 5738-5741; H.W. WHITLOCK et al.: "Behavior of di- and tetrahydro-porphyrins under Alkaline conditions. Direct Observation of the Chlorin-Phlorin Equilibrium" * page 5739, figure 1 *	1,2	
A	--- EP - A - 0127797 (F. HOFFMANN - LA ROCHE AND CO AG) * abstract *	1,2	
A	--- PATENT ABSTRACTS OF JAPAN, vol. 4, no. 190 (C-37) (672), 26th December 1980; & JP - A- 55 130978 (Y. MAT-SUSHIMA) 11.10.1980 ---	1	
			TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
			A61K31/00 C07D487/00
INCOMPLETE SEARCH <p>The Search Division considers that the present European patent application does not comply with the provisions of the European Patent Convention to such an extent that it is not possible to carry out a meaningful search into the state of the art on the basis of some of the claims.</p> <p>Claims searched completely: 1-5: All claims comprise, inter alia, methods for treatment of the human or animal body by surgery or therapy or diagnostic methods (art. 52(4) EPC)</p> <p>Claims searched incompletely:</p> <p>Claims not searched:</p> <p>Reason for the limitation of the search:</p>			
Place of search Berlin		Date of completion of the search 05.06.89	Examiner HASS, C.V.F.
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	



DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl. 7)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
D, A	JOURNAL OF THE AMERICAN CHEMICAL SOCIETY, vol. 91, no. 26 17th December 1969, pages 7485-7489; H. W. WHITLOCK et al.: "Diimide Reduction of Porphyrins" ---		
			TECHNICAL FIELDS SEARCHED (Int. Cl. 7)

THIS PAGE BLANK (USPTO)